




NATURAL LIPOLYSIS

J Eur Acad Dermatol Venereol. 2008, 22, 1465-1470



USE OF MICRODIALYSIS TECHNIQUE TO ASSESS LIPOLYTIC RESPONSIVENESS OF FEMORAL ADIPOSE TISSUE AFTER Endermologie®.

LAFONTAN M.



www.lpgsystems.com

www.cosire-lpg.com

Contact :

infomed@lpgsystems.com

ORIGINAL ARTICLE

Use of the microdialysis technique to assess lipolytic responsiveness of femoral adipose tissue after 12 sessions of mechanical massage technique

C Monteux,*† M Lafontan‡

† LPG Systems Scientific Research, Sophia Antipolis, France

‡ Unité Inserm-UPS 586, IFR-31-Institut Louis Bugnard, Hôpital Rangueil, Toulouse, France

Keywords

cellulite, femoral adipose tissue, microdialysis.

*Corresponding author, LPG Systems – Scientific Research, 2753 Route des Dolines, BP 243 Sophia Antipolis 06905, France, tel. +334 92383900; fax +334 92960965; E-mail: clelia.monteux@pgsystems.com

Received: 21 January 2008,
accepted 29 May 2008

DOI: 10.1111/j.1468-3083.2008.02918.x

Corrections added after online publication on
29 August 2008: The second author's name.

Abstract

Background Adipocytes in femoral areas are known to be metabolically 'silent'. Changes related to fat cell hypertrophy may be involved in the formation of cellulite. A mechanical massage technique, with circulatory and dermatrophic properties, has been shown to have an impact on clinical evaluations (i.e. changes in morphometric measurements) in cellulite areas. Whether this technique affected lipolytic responsiveness in subcutaneous adipose tissue of cellulite areas was not known.

Objective Using a microdialysis technique in subcutaneous adipose tissue, a study was carried out to test the *in situ* incidence of a mechanical massage technique in terms of adipose tissue responsiveness to a lipolytic challenge.

Materials and methods Nine healthy women volunteers with cellulite (grade ≥ 2) were included and treated with 12 sessions of mechanical massage technique (Endermologie®). Microdialysis has been carried out in the femoral adipose tissue in order to assess lipolytic responsiveness via glycerol determination following perfusion of a lipolytic agent (0.1, 1 and 10 μM isoproterenol). Clinical evaluations (measurements of waist, thighs and skin fold) were carried out in parallel. All evaluations were performed before and after treatment.

Results The studied intervention lowered resting dialysate glycerol levels in femoral adipose tissue. The lipid-mobilizing effect of isoproterenol was enhanced after 1 month of treatment. In addition, a clear decrease of morphometric measurements (mean decrease on thighs perimeter: 3.1 to 3.3 cm, $P < 0.01$) was observed.

Conclusion These results suggest an increase in the lipolytic responsiveness of femoral adipose tissue in women with cellulite having undergone 12 sessions of mechanical massage.

Introduction

Lipolysis in human adipocyte is mediated by the sympathetic nervous system activation. Norepinephrine and epinephrine stimulate both the α 2- and β -adrenergic receptors (AR) on the fat cell surface. Differences in the responsiveness of adipose tissue to the adrenergic stimulation are regional and gender related and related to the functional balance between α 2- and β -adrenergic responses.^{1–5} In women, lipid accretion is favoured in the

gluteo-femoral depots,⁶ which are characterized by the abundance of α 2-AR and a relatively sluggish fat turnover.³ Increased expression of α 2-AR – and concomitant decrease of β -AR responsiveness – with fat cell hypertrophy could be a physiological adaptation that leads to the reduction of the lipolytic responsiveness of the hypertrophied adipocyte.⁴ In addition, blood flow was weaker in femoral than in abdominal fat deposits.^{7,8} It remains to establish if the defective lipolytic response and reduced local blood flow (which is supposed to lead to

some fluid stagnation) in those specific fat areas are involved in the aetiology of cellulite.

A mechanical massage technique with circulatory and dermatrophic properties^{9–12} has been shown to have an impact on clinical parameters (i.e. morphometric measurements) in cellulite areas. Whether this technique affected lipolytic responsiveness in subcutaneous adipose tissue (SCAT) of cellulite areas was not known.

Mechanism of lipid mobilization in SCAT can be monitored by microdialysis technique.¹³ It consists of introducing small microdialysis probes in adipose tissue in order to continuously monitor glycerol (marker of lipolysis) in the extracellular space and at the same time deliver drugs locally to this space, thereby eliminating the systemic effects of the drug.¹⁴ This *in vivo* and *in situ* approach allows for the study of adipose cells in their actual milieu and has been largely used. The role of β and α 2-AR in the control of lipolysis in human adipose tissue^{15,16} and the lipolytic responsiveness of adipose tissue in lean and obese subjects¹⁷ have been studied using this technique through measurement of glycerol output combined with estimation of local blood flow changes as it plays an important role for regulating lipid mobilization.¹⁵

The aim of the present study was to explore the lipolysis regulation of subcutaneous femoral adipocytes in female volunteers subjected to 12 sessions of a mechanical massage technique. Isoproterenol-mediated lipolysis and adipose tissue blood flow were assessed, using *in situ* microdialysis, before and after treatment. Clinical parameters (perimetry and skin fold measurements) were also investigated to appreciate morphological changes.

Materials and methods

Subjects

The experimental procedure was approved by the Local Ethics Committee of Toulouse Hospital (France) and was carried out on nine healthy female volunteers who signed an informed consent form and had cellulite of the thighs of grade ≥ 2 according to Nurnberger classification.¹⁸ In order to exclude any interference due to hormonal or environmental effects, the subjects were required to be receiving oral contraception for more than 3 months and to maintain stable dietary habit, weight and physical activity, for the total duration of the study. The characteristics of the subjects are depicted in Table 1.

Experimental design

Before and after treatment, all the evaluations were carried out in the laboratory. Lipolytic response in femoral adipose tissue and local blood flow responsiveness were

Table 1 Characteristics of the subjects (n = 9)

	Mean \pm SE	Range
Age, year	41.2 \pm 3.6	36–45
Height, cm	164.89 \pm 8.34	155–180
Weight, kg	74.37 \pm 13.44	60–97
Body mass index, kg/m ²	27.16 \pm 0.95	22.58–29.94

assessed *in situ* by dosage of glycerol and urea output, respectively, using the microdialysis method. Objective physical measurements included perimetry and skin fold thickness.

Treatment with the mechanical massage technique

The mechanical massage technique (Endermologie®) is a non-invasive technique consisting of a deep tissue mobilization provided by a medical device composed of a treatment chamber with an aspiration system and two independent motorized rollers which roll and unroll a skin fold. Each subject received a total of 12 sessions (thrice a week) of mechanical massage technique according to a specific practical protocol.

Microdialysis experiments

Subjects were investigated at 8:00 a.m. after an overnight fast before and after 12 sessions with the mechanical massage technique. After light epidermal anaesthesia (Emla® patch 5%, AstraZeneca, 20 min), one microdialysis probe (20 \times 0.5 mm, 20-kDa molecular-weight cutoff; CMA Microdialysis, Sweden) was inserted percutaneously into the femoral SCAT at two-thirds of the distance between the patella and the superior anterior iliac spine. The probe was connected to a microinjection pump and continuously perfused with a sterile Ringer solution (154 mM sodium, 6 mM potassium, 2.5 mM calcium, 160 mM chloride). The perfusion was set at a flow rate of 2 μ L/min and fractions were collected following a 30-min equilibration period. After fractions were collected for definition of baseline values, the probe was infused with increasing concentrations of the β -AR agonist isoproterenol (0.1; 1 and 10 μ M, in Ringer solution). Three 10-min fractions were collected for each concentration and frozen (-80 °C) until dosage. Stimulation of lipolysis promotes glycerol appearance in the extracellular fluid which is taken up by the microdialysis probe. Glycerol which is not re-utilized by fat cells (i.e. due to glycerol kinase defect) is the best metabolic index for lipolysis determination.¹³

Biochemical determinations

The concentrations of glycerol and urea in the microdialysis samples were analysed using enzymatic methods as previously described.¹⁹ Urea was used as an endogenous reference compound to determine adipose tissue blood flow.²⁰

Perimetry and cutaneous fold

Perimeter measurements were performed with a millimeter tape measured in one point of each thigh (middle thigh) and at the waist and hips. Caliper gauge measurements of skin fold thickness were performed at one lateral point of the right and left inferior limbs. The height of each measurement point was identified throughout the study using a height gauge.

Statistical analysis

All the values are given as means \pm SE. Wilcoxon's or Student tests for paired values were used for comparison. $P < 0.05$ was considered statistically significant. Concentration-dependent responses to isoproterenol infusions were compared using an ANOVA for repeated measures. All statistical comparisons were performed by using a statistical software package (Stata SE 8.2, Statacorp, College Station, TX).

Results

All patients (nine) completed the 12 sessions. The sample size for analyses was 9, except for the microdialysis measurements where it was 8, due to a technical problem with one patient.

Changes of lipid-mobilizing responses to isoproterenol induced by the mechanical massage technique in femoral adipose tissue

The effect of isoproterenol on glycerol output is depicted in fig. 1.

At basal level (without isoproterenol), glycerol concentration is lowered after treatment but the difference is not statistically significant ($P = 0.0882$).

The addition of isoproterenol into the perfusate increased the dialysate glycerol concentration over the baseline significantly at $1 \mu\text{M}$ before treatment ($P = 0.0108$) and at 0.1 and $1 \mu\text{M}$ after treatment ($P = 0.0038$ and 0.0153 , respectively).

The highest isoproterenol concentration ($10 \mu\text{M}$) did not significantly modify the glycerol output in any condition.




fig. 1 Effect of the non-selective β -AR agonist isoproterenol on the dialysate glycerol concentration in femoral SCAT from women subjects ($n = 8$) before and after treatment (12 sessions).




fig. 2 Effect of the non-selective β -AR agonist isoproterenol on the glycerol evolution, expressed in percentage over the baseline value, in femoral SCAT from women subjects ($n = 8$) before and after treatment (12 sessions).

Results expressed in percentage over the baseline value are shown in fig. 2. After 12 sessions with the mechanical massage technique, responsiveness to isoproterenol infusion is improved for the usually used concentrations in microdialysis experiments on adipose tissue: 50% vs. 27% at $0.1 \mu\text{M}$, 83% vs. 63% at $1 \mu\text{M}$ (ANOVA $P = 0.0441$) and 37.5% vs. 0% at $10 \mu\text{M}$ (ns), respectively, for responses obtained after and before treatment.

Effects of the mechanical massage technique on femoral adipose tissue blood flow to isoproterenol

The effect of isoproterenol on urea concentration is depicted in fig. 3.

At basal level (without isoproterenol), urea concentration is similar before and after treatment (0.82 ± 0.47 vs. 0.78 ± 0.39 mmol/L, ns).




fig. 3 Effect of the non-selective β -AR agonist isoproterenol on urea concentration (endogenous reference compound) in femoral SCAT from women subjects ($n = 8$) before and after treatment (12 sessions).

The addition of isoproterenol into the perfusate did not significantly modify the urea concentration when compared with baseline except at the highest concentration used (10 μM) before treatment and at 1 μM after treatment where it is lowered ($P = 0.004$ and $P = 0.0102$, respectively).

When results are expressed in percentage below the baseline, isoproterenol infusion caused a dose-dependent decrease in urea concentrations from 8.5% at 0.1 μM ($P = 0.2084$) to 18.3% at 1 μM ($P = 0.1451$) and 46.3% at 10 μM ($P = 0.004$) before treatment. The decrease in the urea concentrations promoted by 0.1 and 1 μM isoproterenol is greater after treatment (23% and 25%, respectively), but it is not statistically significant ($P = 0.111$ and $P = 0.257$, respectively).

Effects of the mechanical massage technique on clinical evaluations

Statistical comparison of morphologic parameters are summarized in Table 2. On the whole, the measurements showed a significant reduction after treatment. This reduction is observed on cutaneous fold (−12.89% on the Right side and −13.41% on the Left side), waist and thigh measurements (1.7 to 3.3 cm; $P < 0.01$).

Discussion

The main finding of this study is that treatment of femoral fat deposits with 12 sessions of mechanical massage promoted an increase β -AR-mediated responsiveness in a tissue known to be poorly responsive to catecholamines

Table 2 Circumference and cutaneous fold measurements before and after treatment (12 sessions)

	Before treatment	After treatment	P
Waist measurement (cm)	97.8 \pm 3.6	96.1 \pm 3.6	< 0.01
Hip measurement (cm)	108.7 \pm 4.0	108.6 \pm 4.0	NS
Right thigh measurement (cm)	64.7 \pm 3.8	61.4 \pm 3.0	< 0.01
Left thigh measurement (cm)	64.7 \pm 3.8	61.6 \pm 3.0	< 0.01
Right cutaneous fold (mm)	28.4 \pm 1.7	24.8 \pm 1.4	< 0.001
Left cutaneous fold (mm)	28.4 \pm 1.7	24.6 \pm 1.4	< 0.001

Data are expressed as means \pm SEM. P values were calculated using the Student's t -test ($n = 9$).

and metabolically 'silent' in terms of non-esterified fatty acid release.⁸ Before treatment, the β -AR agonist at 1 μM promoted a weak but significant stimulation of glycerol output. It is the response expected in a tissue known to be poorly responsive to lipolytic agents.⁸ It must be noted that the highest pharmacologic concentration of isoproterenol (e.g. without physiological relevance) is without effect. After treatment, isoproterenol responsiveness was improved as shown by a shift to the left of the concentration–response curve, 0.1 μM isoproterenol becoming efficient. Moreover, percent increase over basal values was enhanced after treatment. Although they had no quantitative relevance, urea results suggest that blood flow changes also occur after treatment with the mechanical massage. However, interpretations are not easy in the absence of further investigations with another blood flow tracer. The limited number of patients involved in the present study allows a clear demonstration of the metabolic impact of this mechanical massage. Nevertheless, it was not possible to correlate isoproterenol-induced lipolytic responses of adipose tissue and the morphologic changes; such a study, which requires a large number of patients, was considered to be out of the scope of the present investigation.

The biphasic concentration response-curve depicted in the figures, when infusing isoproterenol, is difficult to interpret in the absence of more detailed studies of the composition of dialysate. It must be noted that this kind of biphasic curve is obtained when treating isolated fat cells with isoproterenol or norepinephrine *in vitro*.²¹ Sustained stimulation of lipolysis by pharmacological doses of lipolytic agents induces ATP depletion and reduces lipolytic responses. Moreover, since the effects of isoproterenol are performed with sequential and cumulative procedure, it is not excluded that a partial desensitization of β -AR,¹ leading to a reduction of the response, could occur under our working conditions. When performing *in vivo* studies,

it is also possible that at the highest concentration, isoproterenol treatment alters the normal function of adipocytes and endothelial cells. It is possible that infusion of the higher doses strongly alters adenosine and prostaglandin release, two agents known to exert anti-lipolytic effects and possess vascular actions. All these factors could alter glycerol output from the microdialysis probe.

Cellulite is a condition that remains an issue of cosmetic concern to a great number of women. A large part of the difficulties in treating cellulite arises from our incomplete understanding of this phenomenon.^{22–24} Questions concerning aetiology and even definition of cellulite remain largely open and debated. Due to the lack of convincing experimental studies, there is no consensus.

A number of products (herbal extracts, retinol, caffeine, ruscogenin, etc.) with lipolytic, venotonic or trophic properties, are used in numerous slimming treatments (topics, dietary complements, mesotherapy).^{25–29} Modern technological advances using laser or electromagnetic energies claim to improve cellulite,^{30,31} but data are mainly subjective. Finally, efficacy and safety of anti-cellulite treatments described in literature are more or less convincing.²²

The present investigation is the first attempt to characterize the putative functional impact of a treatment with a mechanical massage technique on the responsiveness of the adipose tissue located in cellulite areas. For that purpose, using the microdialysis technique, the lipolytic responses induced by a β -AR agonist in femoral adipose tissue was investigated before and after treatment in women with cellulite. Similar findings (i.e. improvement of the response promoted by isoproterenol in SCAT) were observed in previous studies carried out in abdominal area of obese subjects submitted to physical exercise^{32–35} or hypocaloric diets.^{36,37} Benefits of exercise training have been explored after long training periods. For example, a 4-month endurance training program, improves ANP- as well as β -AR-mediated lipid mobilization and adipose tissue blood flow in the abdominal SCAT of overweight subjects. The recovery of a higher lipolytic efficiency in adipose tissue is an important benefit of a training program in overweight subjects.³⁴ In front of the results obtained after longer periods of endurance training, it is not excluded that longer sessions or longer periods of treatment with the mechanical massage could be more beneficial for a better recovery of lipolytic responsiveness in femoral fat deposits.

In the absence of further investigations and notably follow-up experiments, it is difficult to speculate on the duration of the improvement observed in the lipolytic responsiveness of femoral adipose tissue. Regarding morphological changes in cellulite areas, a remanent effect has been evaluated at 6 months with perimetry,

skin imprints and 20-MHz sonography.¹¹ Nevertheless, based on results of clinical studies in patients performing exercise,^{34,35} the improvement observed in the present study is expected to remain efficient in the long term if and only if the patients maintain some physical exercise and nutritional control.

Conclusion

This study shows that a treatment (12 sessions) with this mechanical massage technique improves the lipid-mobilizing effects of *in situ* administered isoproterenol in femoral adipose tissue of women with cellulite. In addition to these functional improvements in femoral adipose tissue biology, the treatment reduced body circumferences and skin fold measurements. The mechanism of action of such treatment is still unclear. It may initiate local changes (i.e. adipose tissue remodelling events) which have an incidence on the lipolytic responsiveness of femoral adipose tissue.

Complementary studies are necessary to confirm the effect in other fat deposits and to enter more deeply into further mechanistic consideration on the incidence of this mechanical massage technique on lipid mobilization in fat deposits.

Acknowledgements

This study was carried out by Groupe Santé Recherche, 35, rue Bernard de Ventadour, 31300 Toulouse (France) and was funded by LPG Systems, the makers of the medical device used in this study.

References

- Ostman J, Arner P, Engfeldt P, Kager L. Regional differences in the control of lipolysis in human adipose tissue. *Metabolism* 1979; **28**: 1198–1205.
- Williams CM. Lipid metabolism in women. *Proc Nutr Soc* 2004; **63**: 153–160.
- Frayn KN. Regulation of fatty acid delivery *in vivo*. *Adv Exp Med Biol* 1998; **441**: 171–179.
- Lafontan M, Berlan M. Do regional differences in adipocyte biology provide new pathophysiological insights? *Trends Pharmacol Sci* 2003; **24**: 276–283.
- Bjorntorp P. The regulation of adipose tissue distribution in humans. *Int J Obes Relat Metab Disord* 1996; **20**: 291–302.
- Rebuffe-Scrive M, Enk L, Crona N *et al*. Fat cell metabolism in different regions in women. Effect of menstrual cycle, pregnancy, and lactation. *J Clin Invest* 1985; **75**: 1973–1976.
- Millet L, Barbe P, Lafontan M, Berlan M, Galitzky J. Catecholamine effects on lipolysis and blood flow in human abdominal and femoral adipose tissue. *J Appl Physiol* 1998; **85**: 181–188.

- 8 Tan GD, Goossens GH, Humphreys SM, Vidal H, Karpe F. Upper and lower body adipose tissue function: a direct comparison of fat mobilization in humans. *Obes Res* 2004; **12**: 114–118.
- 9 Chang P, Wiseman J, Jacoby T, Salisbury AV, Ersek RA. Noninvasive mechanical body contouring: Endermologie a one-year clinical outcome study update *Aesthetic Plast Surg* 1998; **22**: 145–153.
- 10 Innocenzi D, Balzani A, Montesi G *et al.* Evidenza delle modificazioni cutanee indotte dalla Tecnica LPG® mediante analisi d'immagine. *Derma Cosmetologia* 2003; **2**(1): 9–15.
- 11 Ortonne JP, Queille-Roussel C, Duteil L *et al.* Treatment of cellulite: effectiveness and sustained effect at 6 months with Endermologie demonstrated by several quantitative evaluation methods. *Nouv Dermatol* 2004; **23**: 261–269.
- 12 Moseley A, Esplin M, Piller N, Douglas J. Endermologie: a new treatment option for secondary arm lymphedema. *Lymphology*. 2007; **40**: 129–137.
- 13 Lafontan M, Arner P. Application of *in situ* microdialysis to measure metabolic and vascular responses in adipose tissue. *Trends Pharmacol Sci* 1996; **17**: 309–313.
- 14 Arner P, Bolinder J, Eliasson A, Lundin A, Ungerstedt U. Microdialysis of adipose tissue and blood for *in vivo* lipolysis studies. *Am J Physiol* 1988; **255**: E737–E742.
- 15 Galitzky J, Lafontan M, Nordenstrom J, Arner P. Role of vascular alpha-2 adrenoceptors in regulating lipid mobilization from human adipose tissue. *J Clin Invest* 1993; **91**: 1997–2003.
- 16 Barbe P, Millet L, Galitzky J, Lafontan M, Berlan M. *In situ* assessment of the role of the beta 1-, beta 2- and beta 3-adrenoceptors in the control of lipolysis and nutritive blood flow in human subcutaneous adipose tissue. *Br J Pharmacol* 1996; **117**: 907–913.
- 17 Jansson PA, Larsson A, Smith U, Lonnroth P. Glycerol production in subcutaneous adipose tissue in lean and obese humans. *J Clin Invest* 1992; **89**: 1610–1617.
- 18 Nurnberger F, Mullr G. So-called cellulite: an invented disease. *J Dermatol Surg Oncol* 1978; **4**: 221–229.
- 19 Belhadj-Tahar H, Nassar BB, Sadeg N, Coulais Y. Monitoring of adipocyte responsiveness by *in situ* microdialysis in lipodystrophy tissue: adjustment of a glycerol quantification method in small samples. *Acta Clin Belg* 2006; **61**: 143–146.
- 20 Schwalbe O, Buerger C, Plock N, Joukhadar C, Kloft C. Urea as an endogenous surrogate in human microdialysis to determine relative recovery of drugs: analytics and applications. *J Pharm Biomed Anal* 2006; **41**: 233–239. Epub 2005 December 20.
- 21 Stich V, Harant I, deGlisezinski I *et al.* Adipose tissue lipolysis and hormone-sensitive lipase expression during very-low-calorie diet in obese female identical twins. *J Clin Endocrinol Metab* 1997; **82**: 739–744.
- 22 Avram MM. Cellulite: a review of its physiology and treatment. *J Cosmet Laser Ther* 2004; **6**: 181–185.
- 23 Callaghan T, Wilhelm KP. An examination of non-invasive techniques in the analysis and review of cellulite. *J Cosmet Sci* 2005; **56**: 379–393.
- 24 Draelos ZD. The disease of cellulite. *J Cosmet Dermatol* 2005; **4**: 221–222.
- 25 Hexsel D, Orlandi C, Zechmeister do Prado D. Botanical extracts used in the treatment of cellulite. *Dermatol Surg* 2005; **31**: 866–872.
- 26 Pierard-Franchimont C, Pierard GE, Henry F, Vroome V, Cauwenbergh G. A randomized, placebo-controlled trial of topical retinol in the treatment of cellulite. *Am J Clin Dermatol* 2000; **1**: 369–374.
- 27 Soler-Rouanet B, Maille PA, Adhoute H, Dubertret L. Etude par photogrammétrie de l'efficacité amincissante du complément alimentaire «Cellulase Urto» chez des volontaires sains féminins. *Nouv Dermatol* 2001; **20**: 13–18.
- 28 Bertin C, Zunino H, Pittet JC *et al.* A double-blind evaluation of the activity of an anti-cellulite product containing retinol, caffeine, and ruscogenine by a combination of several non-invasive methods. *J Cosmet Sci* 2001; **52**: 199–210.
- 29 Rotunda AM, Avram MM, Avram AS. Cellulite: Is there a role for injectables? *J Cosmet Laser Ther* 2005; **7**: 147–154.
- 30 Sadick N, Magro C. A study evaluating the safety and efficacy of the VelaSmooth system in the treatment of cellulite. *J Cosmet Laser Ther* 2007; **9**: 15–20.
- 31 Nootheti PK, Magpantay A, Yosowitz G, Calderon S, Goldman MP. A single center, randomized, comparative, prospective clinical study to determine the efficacy of the Velasmooth system versus the Triactive system for the treatment of cellulite. *Lasers Surg Med* 2006; **38**: 908–912.
- 32 Arner P, Kriegholm E, Engfeldt P, Bolinder J. Adrenergic regulation of lipolysis *in situ* at rest and during exercise. *J Clin Invest* 1990; **85**: 893–898.
- 33 Stich V, de Glisezinski I, Galitzky J *et al.* Endurance training increases the beta-adrenergic lipolytic response in subcutaneous adipose tissue in obese subjects. *Int J Obes Relat Metab Disord* 1999; **23**: 374–381.
- 34 Moro C, Pillard F, De Glisezinski I *et al.* Training enhances ANP lipid-mobilizing action in adipose tissue of overweight men. *Med Sci Sports Exerc* 2005; **37**: 1126–1132.
- 35 De Glisezinski I, Crampes F, Harant I *et al.* Endurance training changes in lipolytic responsiveness of obese adipose tissue. *Am J Physiol* 1998; **275**: E951–E956.
- 36 Barbe P, Stich V, Galitzky J *et al.* *In vivo* increase in beta-adrenergic lipolytic response in subcutaneous adipose tissue of obese subjects, submitted to a hypocaloric diet. *J Clin Endocrinol Metab* 1997; **82**: 63–69.
- 37 Sengenès C, Stich V, Berlan M *et al.* Increased lipolysis in adipose tissue and lipid mobilization to natriuretic peptides during low-calorie diet in obese women. *Int J Obes Relat Metab Disord* 2002; **26**: 24–32.